

POST OPERATIVE CARE BREAST LIFT (MASTOPEXY)

Post op care:

- Surgical bra to be worn 24/7 until follow up appointment in 7-10 days
- Sleep on back, or side if comfortable, with head, neck, and shoulders elevated
- May shower 24 hours after surgery. Leave white tapes (steri-strips) on skin. These may get wet. They may fall off after a couple days.
- No straining, lifting, or bending for 2 weeks
- Ok to take Tylenol immediately and Ibuprofen immediately after surgery (do not take with other acetaminophen medications such as Percocet)
- Can operate a motor vehicle 24 hours after last dose of narcotic pain medication
- Rest – can resume most activities in 2 weeks and exercise in 4 weeks
- Scopolamine patches can cause temporary pupil dilation and blurry vision. This is usually not serious. Remove the patch immediately and contact our office.

What to expect:

- Swelling and bruising can occur up to 48 hours post-op. May take several weeks (bruising) to several months (swelling) to resolve
- Moderate discomfort – take pain medication as directed
- Constipation may occur with narcotic pain medicine – increase fluid intake and use mild laxative or stool softener if needed
- Asymmetry – right and left sides will heal at different rates
- Numbness, tingling, and sensitivity to physical contact may persist for several weeks or longer

Follow-up:

- Follow up appointment 7-10 days post operatively
- Bring sports bra to follow up appointment unless bra is being provided by this office
- May return to work 1 week post op unless otherwise directed
- No underwire bra in bathing suit for 3 months

**DO NOT HESITATE TO CALL THE OFFICE WITH ANY QUESTIONS
OR CONCERNS**

Pain medication: _____

Antibiotic: _____

AESTHETIC SURGERY
Rosemont Business Campus
919 Conestoga Rd.
Bldg 1, Suite 200
Rosemont, Pa. 19010

RONALD A. LOHNER, M.D., F.A.C.S.

RECONSTRUCTIVE SURGERY

Phone: 610-519-0600

Fax: 610-519-1238

Follow-up: _____